

## Records of coprophilous fungi – a data set

MICHAEL J. RICHARDSON\*

165 Braid Road, Edinburgh, UK EH10 6JE

\* CORRESPONDENCE TO: [mjrichardson@clara.net](mailto:mjrichardson@clara.net)

**ABSTRACT** — Many coprophilous fungi are adapted to their habitat by having specialised spore dispersal mechanisms to improve the chance of the spores being consumed, since they require passage through an animal's gut to encourage their germination. Analysis of coprophilous fungus records allows a study of regional, substrate, and seasonal differences, and has demonstrated that fungi increase in biodiversity with decreasing latitude. 1386 samples of dung, mainly of herbivorous mammals and birds, were collected, mostly between 1994 and 2014, from the wild from various parts of the world. Most (88%) were from north temperate areas, with 7% from south temperate areas, 4% from the tropics, and 0.3% from the Arctic. Over 12,300 records of coprophilous fungi were obtained from these samples on incubation in damp chambers. Details of the fungi found and their origin are presented as files that provide a data set that triples the amount of data used in a 2001 analysis. The data set includes collection details for the samples (locality, country, latitude/longitude coordinates, elevation), date of collection, dung type, species recorded from each sample, location of herbarium deposits, and citations to references where the records have been published. Notes of observations made on each sample during incubation [one pdf for each sample] are also in the on-line checklist, with pdfs of publications in which the records are cited.

**KEY WORDS** — distribution, diversity, dung fungi, substrate preference, worldwide

### Introduction

Coprophilous fungi are suitable subjects for many studies of distribution and diversity; their substrate is abundant and widespread, and readily collectable for drying and return to the laboratory for rehydration and incubation in damp chambers. They are adapted to their habitat by having specialised spore dispersal mechanisms to improve the chance of the spores being consumed, since they require passage through an animal's gut to encourage their germination. These adaptations include positive phototropism of sporophores; ballistic spore discharge around midday to better get spores into the airstream; gelatinous appendages to allow them to adhere to vegetation; and spore pigmentation to protect from UV damage while exposed. Analysis of coprophilous fungus records have allowed regional, substrate and seasonal differences in the occurrence of this specialist group of fungi to be studied, and have demonstrated that the increase in biodiversity with decreasing latitude, well known for other organisms, also applies to fungi (Richardson 2001). 1386 samples of dung, mainly of herbivorous mammals and birds, were collected, mostly between 1994 and 2014, from the wild from various parts of the world. Caution is needed when interpreting the results of limited collections made *en passant*, since it is well known that observations from such collections cannot replace an intensive and comprehensive study of collections made over a period of time from a locality. Similarly, interactions among competing fungi and other biota could well influence which fungi actually fruit when samples are incubated. It is hoped,

however, that the records from these collections will contribute to our knowledge of the distribution and occurrence of coprophilous fungi. The now relatively aged monographs of many of the common coprophilous genera, while still invaluable, are much in need of revision and updating to facilitate these studies. A summary of the geographical origin of the samples that provided the records is given in Table 1. Over 12300 records of coprophilous fungi were obtained from these samples on incubation in damp chambers. Details of the fungi found and their origin are presented as files in the on-line version of this paper. The files provide a data set that triples the amount of data used in the earlier analysis (Richardson 2001). The data set includes collection details for the samples (locality, country, latitude/longitude coordinates, elevation), date of collection, dung type, species recorded from each sample, location of herbarium deposits and citations to references where the records have been published. Notes of observations made on each sample during incubation [one pdf for each sample] are also in the on-line version, with pdfs of publications in which the records are cited.

TABLE 1. Summary of data

STATES	NO. OF SAMPLES	NO. OF RECORDS	NO. OF SPECIES (APPROX.)*
Australia	46	545	146
British Overseas Territory <sup>†</sup>	9	72	39
Brazil	7	75	32
Canada	11	50	33
Chile	2	14	11
Costa Rica	3	26	20
Dominica	4	40	28
Egypt	1	7	7
Falkland Islands	39	215	73
Faeroe Islands	20	233	59
Finland	30	199	78
France <sup>§</sup>	101	978	182
Great Britain [inc. Isle of Man]	831	7582	317
Greece	43	344	79
Greenland	1	1 <sup>z</sup>	0
Iceland	76	671	115
Eire	5	44	31
Italy	11	133	75
Malaysia	1	16	16
Morocco	14	167	57
Netherlands Antilles	12	98	37
New Zealand	1	1	1
Norway	3	24	20
Papua New Guinea	2	4	4
Puerto Rico	5	45	26
Spain	24	221	93
St Lucia	5	52	29
Sweden	19	110	37
Tunisia	4	21	16
US Virgin Islands	2	16	14
USA	51	319	128
<b>TOTAL</b>	<b>1383</b>	<b>12323</b>	<b>566</b>

\* excluding potential duplicates of identifications such as cf., aff., sp., ?, etc., to prevent double counting.

† Ascension Island and St Helena. § including Corsica, Guadeloupe and the Kerguelen Islands.

<sup>z</sup> null record from the single Greenland sample.

**Acknowledgments**

Many thanks to seventy friends and colleagues who have collected samples for me from around the world. Thanks also to Ann Bell, Francesco Doveri and Roy Watling for acting as expert reviewers, to Noni Korf for facilitating the on-line presentation, and to Lorelei Norvell for advice and considering the publication of a data paper in *Mycotaxon*.

**Literature cited**

Richardson MJ. 2001. Diversity and occurrence of coprophilous fungi. *Mycological Research* 105: 387-402. <http://dx.doi.org/10.1017/S0953756201003884>

**Literature cited for records in the dataset**

All those identified by \* are available in the on-line files of this paper

- Bramley WG. 1985. *Fungus Flora of Yorkshire 1985*. Yorkshire Naturalists' Union, Leeds. 277pp.
- \*Huhtinen S, Spooner BM. 2003. A redistribution of *Ascophanus tityrii* Velen. *Kew Bulletin* 58: 749-757. <http://dx.doi.org/10.2307/4111156>
- Krug JC, Cain RF. 1974. New species of *Hypocopra* (*Xylariaceae*). *Canadian Journal of Botany* 52: 809-843. <http://dx.doi.org/10.1139/b74-105>
- Richardson MJ. 1968. Local Society Forays. *Bull. Brit. Mycol. Soc.* 1968: 2:17
- Richardson MJ. 1970. Report of the cryptogamic section 1966. *Trans. Bot. Soc. Edinburgh* 40: 350-352. <http://dx.doi.org/10.1080/03746606708685159>
- Richardson MJ. 1970. Report of the cryptogamic section 1967. *Trans. Bot. Soc. Edinburgh* 40: 471-472.
- Richardson MJ. 1970. Report of the cryptogamic section 1968. *Trans. Bot. Soc. Edinburgh* 40: 648-650. <http://dx.doi.org/10.1080/03746606908685189>
- \*Richardson MJ. 1970. *Ballocephala verrucospora* sp. nov. parasitizing Tardigrades. *Transactions of the British Mycological Society* 55: 307-309.
- \*Richardson MJ. 1972. Coprophilous ascomycetes on different dung types. *Transactions of the British Mycological Society* 58: 37-48. [http://dx.doi.org/10.1016/S0007-1536\(72\)80069-X](http://dx.doi.org/10.1016/S0007-1536(72)80069-X)
- \*Richardson MJ. 1998. New and interesting records of coprophilous fungi. *Botanical Journal of Scotland* 50: 161-175. <http://dx.doi.org/10.1080/03746609808684913>
- \*Richardson MJ. 1998. *Coniochaeta polymegasperma* and *Delitschia trichodelitschioides*, two new coprophilous ascomycetes. *Mycological Research* 102: 1038-1040. <http://dx.doi.org/10.1017/S0953756297005558>
- \*Richardson, M.J. (2001). Diversity and occurrence of coprophilous fungi. *Mycological Research* 105: 387-402. <http://dx.doi.org/10.1017/S0953756201003884>
- \*Richardson MJ. 2001. Coprophilous fungi from Brazil. *Brazilian Archives of Biology and Technology* 44: 283-389. <http://dx.doi.org/10.1590/S1516-89132001000300010>
- \*Richardson MJ. 2004. Coprophilous fungi from Iceland. *Acta Botanica Islandica* 14: 77-103.
- \*Richardson MJ. 2005. Coprophilous fungi from The Faroe Islands. *Fróðskaparrit* 53: 67-81.
- \*Richardson MJ. 2005. Moroccan coprophilous fungi. *Botanical Journal of Scotland* 56: 147-162. <http://dx.doi.org/10.1080/03746600408685075>
- \*Richardson MJ. 2006. New records of fungi from Orkney and Shetland. *Botanical Journal of Scotland* 58: 93-104. <http://dx.doi.org/10.1080/03746600608685111>
- \*Richardson MJ. 2006. A new species of *Coniochaeta* from Perthshire. *Botanical Journal of Scotland* 58: 105-107. <http://dx.doi.org/10.1080/03746600608685112>
- \*Richardson MJ. 2007. The distribution and occurrence of coprophilous *Ascobolaceae*. *Mycologia Montenegrena* 10: 211-227.
- \*Richardson MJ. 2008. Records of French coprophilous fungi. *Cryptogamie, Mycologie* 29: 157-177.
- \*Richardson MJ. 2008. Coprophilous fungi from the Greek Aegean islands. *Mycologia Balcanica* 5: 23-32.
- \*Richardson MJ. 2008. Records of Coprophilous Fungi from the Lesser Antilles and Puerto Rico. *Caribbean Journal of Science* 44: 206-214.

- \*Richardson MJ. 2011. Additions to the coprophilous mycota of Iceland. *Acta Botanica Islandica* 15: 23-50.
- \*van Brummelen J, Richardson MJ. 2000. *Ascozonus monascus*, a new species of ascomycetes from Great Britain. *Persoonia* 17: 487-490.
- \*Watling R, Richardson MJ. 2010. Coprophilous fungi of the Falkland Islands. *Edinburgh Journal of Botany* 67: 399-423. <http://dx.doi.org/10.1017/S0960428610000156>
- \*Webster J, Whalley AJS, Thienhirung Surang, Richardson MJ. 1999. *Wawelia argentea* and *W. microspora*, two new species of xerophilous fungi on rabbit and hare dung in Britain. *Mycological Research* 103: 1604-1608. doi:10.1017/S0953756299008837