

Horse Banito Tlato Grass do SOT Brazil 6.7.98 58/98
56° 30' 2" W 21° 0' 30" S incubated 31.7

3.8 - 1.8 -

10.8

19.8 Saccobus yellow 4 x 2

swiss 48 x 16

Spores smooth 19.5 x 6.5

Citrus (2)

Coprinus cordisporus ✓ 4-sp

6.5 x 6.5 x 4

(7)

Spiky Podospore 4 sp. 32-34

Panzeria ✓ (4)

24.8

Podospore 8-spaced 32-35.3 x ~~19.3~~

bracket

ly app 20 hrs.

2 hrs

4 at top of ly at 4 at other end

Common ✓ (3)

White Coprinus 9-10 x 6.5-7 ellipsoid

8 sp. Ascii 80-100 x 16 1-2 seriate

long clavate

Paras non-capitate, hyaline
lactose ? (1)

25.8

Coprinus Ascii 80 x 14

8 sp hyaline ellipsoid

Spores 11-11.5 x 6.5-7

Pod. commonis

Pod. imitate with very broad beaded blunt asci

Saccobolus 4 x 2 sm 48 x 16 Citrus ✓
v st. rough. yellow pores

Asci 120 x 38

Sp. minima spores fragmented 30-32 x 5 (5)

5.9. Podospira 8 sp Common ✓

353 x 18

Copros pure white spores 8-10 x 6.5
asci 75-80 x 16 Lactaria

11.9 Cop. lactaria Copros lactaria (8)

Zytopore zygospor. (6) intercalary cell. 9u long
dark cells v. variable 22.5 x 10 to 42 x 16

17.9

Imm Copros? copros spores on old
stipes 6 x 6 x 4

P. Complanata

~~Discarded~~ 17.9

Discarded

- not much on it +
rather mites / nearby